



INTERNATIONAL JOURNAL OF ADVANCE RESEARCH, IDEAS AND INNOVATIONS IN TECHNOLOGY

ISSN: 2454-132X

Impact factor: 4.295

(Volume3, Issue1)

Bioprospecting of Hibiscus Rosasinensis for Antimicrobial Activity against Soil Microbes

R. Prasanna

PRIST University, Tamilnadu

prasanna16.bt@gmail.com

Abstract: The present investigation was carried out about the evaluation of antimicrobial activity against corrosion causing soil microbes on buried elements in acidified mediated soil by Agar disc diffusion method using Aqueous and Ethanol extract of Hibiscus rosasinensis leaves. This plant was subjected to solvent extraction with water and ethanol on increasing the polarity to identify and isolate the antimicrobial active materials. Crude aqueous and ethanol extract of Hibiscus is significantly inhibited the important microbes which causing corrosion such as *S.aureus*, *Streptococcus*, *B. subtilis*, *Lactobacillus*, *Proteus*, *Cornybacterium*, *Pseudomonas*, *A.niger*, *Mucor* and *Desulphovibro sp.* Results of the present studies investigation suggests that Hibiscus are important effective plant for further work on isolating and characterizing of antimicrobial active materials.

Key words: Bioprospecting, *Hibiscus rosasinensis*, *S.aureus*, *Streptococcus*, *B. subtilis*, *Lactobacillus*, *Proteus*, *Cornybacterium*, *Pseudomonas*, *A.niger*, *Mucor* and *Desulphovibro sp.*, Agar well diffusion.

I. INTRODUCTION

Soil is the region for the proliferation of microbes with many colonies which developing around soil particles. Many microbes in soil habitats normally are much higher than those in water. Microbes in soil are important because they affect soil structure and fertility. Soil contains microbes which involves the corrosion of metal which is under the soil in the form of electric cables, sewage and drainage pipe system in acidic mediated soil. Medicinal plants are rich sources of antimicrobial agents.

In present studies, Hibiscus rosasinensis leaves extract used to control the corrosion causing microbes present in acidic mediated soil. Considering the vast potentiality of plants as sources for antimicrobial reagents, a systematic study was undertaken to screen the local flora for antimicrobial activity from Hibiscus leaves. The inhibition of microbial growth in vitro by the extracts of Hibiscus leaves could be due to the presence of some bioactive compounds in the extracts.

Most of microbes such as sulphate or chloride reducing microbes are anaerobic in condition, with pH=6.8 that contain sulphate and chloride, organic substances and some minerals which are live in poorly drained out at 20°C - 30°C. During metabolism of microbes, the extracted oxygen from sulphate and chloride converts soluble and insoluble substances which are easily attacked the metal surfaces. The use of plant extract not only for medicinal uses but also for inhibitory property of corrosion is the most important. Hence the present study was conducted to express the inhibitory activity of selected plants against corrosion caused by soil microbes when extraction of leaves of both aqueous and ethanol solvents.

II. MATERIALS AND METHODS

A. SOIL SAMPLE COLLECTIONS

The soil samples used for this work were collected from 3 different locations near Krishnarajapura area in Bangalore. The samples were labeled according to the site of collection as Garden soil samples, Sewage soil samples and Household soil samples. The samples were transported in polyethylene bags in ice pack to the laboratory. When samples could not process immediately, they were stored at 4°C for no longer than 18 to 24 h.

B. MICROBIOLOGICAL ANALYSES

The soil sample was mixed, and a suspension of 1 g (dry weight equivalent) in 10 ml of sterile water was prepared. 1 ml of the soil suspension was then diluted serially (ten-fold) and used in the estimation of aerobic heterotrophic bacterial and fungal populations by standard spread-plate dilution method described by Seeley and VanDemark (1981), in triplicate. Nutrient agar containing 0.015% (w/v) nystatin (to inhibit fungi growth) was used for bacteria isolation and incubation was at 35°C for five days. Potato dextrose agar to which 0.05% (w/v) chloramphenicol has been added (to inhibit bacteria growth) was used for fungal isolation, and incubation was at ambient temperature for seven days. Pure isolates of representative communities were maintained on agar slant at 4°C. Identification of isolates was based on cultural, microscopic, and biochemical characteristics with reference to Bergey's manual of determinative bacteriology (1989) for bacteria, and Talbot (1978) for fungi.

C. DETERMINATION OF SOIL PH

Soil pH was determined according to the procedure described by Akpor et al. (2006) using Horiba make D-51 pH meter - Measuring object & amp.

D. BIOLOGICAL SAMPLE COLLECTIONS

Leaves of the *Hibiscus rosasinensis* were collected from in the Krishnarajapura area, Bangalore, Karnataka. It was ensure that the plant was healthy and uninfected. The leaves were washed under running tap water to eliminate dust and other foreign particles and to clean them thoroughly and a particular amount of leaves dried under shadow and some fresh leaves kept.

E. PREPARATION OF AQUEOUS EXTRACT OF PLANT SAMPLES

10 grams of air dried powder *Hibiscus rosasinensis* leaves were placed in different distilled water bath and boiled for 6 h. At intervals of 2 h it was filtered through 8 layers of muslin cloth and centrifuged at 5000 x g for 15 min. The supernatant of different samples was collected. After 6 h, the supernatant of different samples was concentrated to make the final volume one-fourth of the original volume. Finally 10 g of each material was extracted in 25 ml of distilled water giving a concentration of 40 mg/0.1 ml. It was then autoclaved at 121 °C and 15 lbs pressure and stored at 4 °C.

F. PREPARATION OF ETHANOL EXTRACT OF PLANT SAMPLES

10 grams of air dried powder *Hibiscus rosasinensis* leaves were placed in 100 ml of organic solvent (ethanol) in a conical flask, plugged with cotton and then kept on a rotary shaker at 190-220 rpm for 24 h. After 24 h, it was filtered through 8 layers of muslin cloth and centrifuged at 5000 x g for 15 min. The supernatant of different samples was collected and the solvent was evaporated to make the final volume one-fourth of the original volume, giving a concentration of 40 mg/0.1 ml. It was stored at 4 °C in airtight bottles for further studies.

G. ASSAY OF ANTIMICROBIAL ACTIVITY USING AGAR WELL DIFFUSION METHOD

The 20 ml of sterilized Muller Hinton Agar for bacteria and Potato dextrose agar for fungal was poured into sterile petriplates, after solidification, 100µl of fresh culture of microbes were swabbed on the respective plates. The wells were punched over the agar plates using sterile gel puncher at concentration 100µl of each plant extract were added to the wells. The plates were incubated for 24 hours at 37°C. After incubation the diameter of inhibitory zones formed around each discs were measured in mm and recorded.

III. RESULT

The antimicrobial activities of *Hibiscus rosasinensis* leaves were carried out. Most of the extract shows an antibacterial and antifungal activity against the human pathogens and also corrosion such as *S.aureus*, *Streptococcus*, *B. subtilis*, *Lactobacillus*, *Proteus*, *Cornybacterium*, *Pseudomonas*, *A.niger*, *Mucor* and *Desulphovibro* sp. Both gram-positive and gram-negative bacteria were sensitive to the extracts.

SOIL pH


The pH values ranged from 6.3 – 7.3. The soil pH in Sewage soil was higher than household soil and garden soil. However, differences in the soil pH values of the different sampling locations were not observed to be statistically significant.

MICROBIAL ANALYSIS OF SOIL

The occurrence of aerobic and anaerobic heterotrophic microbes presents in the different soil samples.

MICROBES	GARDEN SOIL	HOUSEHOLD SOIL	SEWAGE SOIL
AEROBIC BACTERIA			
Lactobacillus sp.	Present	Present	Absent
Bacillus sp.	Present	Present	Present
Proteus	Present	Absent	Absent
Pseudomonas	Present	Present	Present
Micrococcus	Present	Absent	Absent
Streptococcus	Absent	Present	Present
Staphylococcus	Present	Present	Present
Corynebacterium	Present	Present	Present
ANAEROBIC BACTERIA			
Desulphovibrio	Present	Present	Present
D. Africans	Present	Present	Present
FUNGI			
Asperigillus niger	Present	Present	Present
Penicillium	Present	Present	Present
Mucor	Present	Present	Present
Trichophyton	Present	Present	Present
Microsporium	Present	Present	Present

The crude extracts containing multiple organic components including flavonoids, tannins, alkaloids, triterpenoids, all of which are known to have antibacterial affects. Hibiscus leaves extract contain phenolics compounds like tannins that are very good antimicrobial agent. Thus it may be summarized that the class of natural compounds must exhibit the antimicrobial activity. The metabolites have been shown to be responsible for various therapeutic activities of medicinal plants <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3609315/> - b16. Flavonoids especially are known to be effective antimicrobial agent against a wide array of microorganisms. The activity is attributed to their ability to complex with extra cellular and soluble proteins and with bacterial cell wall.



ANTIMICROBIAL ACTIVITY OF AQUEOUS EXTRACT OF HIBISCUS ROSASINENSIS IN AGAR WELL DIFFUSION METHOD (MM).

Disc and well diameter = 6 mm
 Zone of inhibition (clausro et.al)

Interpretative ranges	range (mm)	Descriptive
zone of inhibition	< 10	Resistant
zone of inhibition	10 to 13	Intermediate
zone of inhibition	14 to 19	Sensitive
zone of inhibition	> 19	very sensitive
no zone	----	-----

Thus the plant extracts can be used as an important antibiotic to cure disorders caused by the different strains of microbes. The present studies conclude these extract could inhibit human pathogens growth and also corrosion. The results are encouraging but precise assessment is utterly necessary before being situate in practice as well as the most active extracts can be subjected to isolation of the therapeutic antimicrobials and undergo secondary pharmacological evaluation.

The results obtained shows that different strains responded differently to both aqueous and ethanol fresh *Hibiscus rosasinensis* leaves extracts. On these strains, aqueous and ethanol extracts of fresh *Hibiscus rosasinensis* leaves used shows the zone of inhibition in mm formed. The zone of inhibition for aqueous and ethanol extract decreases as the dilution factor increases. According to graph, ethanol extract is more efficient than aqueous extract of fresh *Hibiscus rosasinensis* leaves. Ethanol extract is more efficient against *Proteus*, *Pseudomonas*, *Cornybacterium*, *B.subtilis* and *Desulphovibrio* species than aqueous extract. Aqueous extract is more efficient against *B.subtilis* and *Lactobacillus* species than ethanol extract. Both aqueous and ethanol extract have equal efficient against the remaining of the strains. No effect of *Hibiscus rosasinensis* against *A.niger*. *B.subtilis*, *Proteus* and *Desulphovibrio* are sulphate and chloride reducing microbes which causes the corrosion on buried metal in acidic mediated soil.

Ethanol extract of fresh *Hibiscus rosasinensis* leaves were more effective. The plant extract contains some phytochemical substances which are secondary metabolites. The active constituents which are present in the leaves are slightly soluble in both aqueous and ethanol solvents such as Terepenoids, tannins, alkaloids, etc. Extract of aqueous and ethanol of fresh *Hibiscus rosasinensis* leaves is compared to consider that ethanol extract is more effective than aqueous. All extracts used during this study, the microbial activity increased with the increase of extract concentration, this means that the zone of inhibition was higher on plates that contain extract with low dilution factor, observed in report done by Esimone et.al. (1998) which says that the plant extracts inhibit the growth of various microbes at different concentration. Like this, similar results recorded obtained where the increase in the concentration of extracts corresponds to the increase of diameter of zone of inhibition. In this study, the active constituents may be present in insufficient quantities in the plant extracts to show activity with the different concentration.

CONCLUSION

Ethanol extracts were more efficient in all cases of fresh *Hibiscus rosasinensis* leaves. This study reveals about antimicrobial effect of *Hibiscus rosasinensis* will make it easier for dosage determination and chemotherapeutic index of the extract. This study informs that antimicrobial effect of *Hibiscus rosasinensis* can change depending on *Hibiscus rosasinensis* plant extract, used, the solvent used, moreover the extract concentration of the fresh *Hibiscus rosasinensis* leaves because each extract has its minimum inhibition concentration (MIC) which is the highest dilution of a plant extract that still retain an inhibitory effect against the microbes growth which causes corrosion. These plant extract inhibits the growth of corrosion causing microbes occur in soil.

REFERENCE

1. Strobel, G., and B. Daisy. 2003. Bioprospecting of microbial endophytes and their natural products. *Microbiol. Molecular. Biol. Rev.* 67(4): 491-502.
2. Drobniowski FA. *Bacillus cereus* and related species. *Clin Microbiology*
3. W. Sand, T. Gehrke, *Research in Microbiology* 157 (2006)
4. Konning GH, Ayare C, Ennison B. Antimicrobial activity of some medicinal plants from Ghana.
5. www.wikipedia.com.
6. Sharma Pankaj et al; Review on neem (azadirachta indica): Thousand
7. problem one solution
8. Kirtikar KR, Basu, BD. In *Medicinal Plants* (eds Blatter, E., Cains, J. F., Mhaskar, K. S.), Vivek Vihar, New Delhi, 1975
9. . Esimone CO, Adiukwu MU, Okonta JM. Preliminary Antimicrobial Screening of the Ethanol Extract from the Lichen *Usnea subfloridans* (L).
10. Seeley and VanDemark (1981)
11. Bergey's manual of determinative bacteriology (1989) for bacteria, and Talbot (1978) for fungi.
12. Akpor et al. (2006)
13. Esimone et.al. (1998)
14. Bergey's manual of determinative bacteriology (1989) for bacteria, and Talbot (1978) for fungi.