

ISSN: 2454-132X

Impact factor: 4.295

(Volume3, Issue3) Available online at www.ijariit.com

# Homomorphism and Isomorphism of Rough Group

L. Jesmalar

Holy Cross College jesmalar2711@gmail.com

Abstract: The chief objective of this study is to show the usefulness of Rough set theory in Group theory. The aim of this paper is to determine the definable and undefinable groups also construct the rough group, upper rough group, homomorphism, and isomorphism of the rough group using rough set theory.

Keywords: Rough Group, Upper Rough Group, Rough Subgroup, Homomorphism and Isomorphism of Upper Rough Group.

# 1. INTRODUCTION

Rough set theory has been combined with other mathematical theories such as modal logic Boolean algebra, fuzzy sets, semigroup, and random set. Among these research aspects, many papers have been focused on the connection between rough sets and algebraic systems. Biswas and Nanda [1, 2] defined the notion of rough subgroups. Kuroki [4] introduced the notion of a rough idea in a semigroup. In my previous papers [8,9,10], rough graph, rough lattice were defined. In this paper, rough group, the upper rough group are defined and explained about the homomorphism and isomorphism of the upper rough group.

# 2. ROUGH SET

# 2.1 Definition [5, 6]

A *rough set* is a formal approximation of a crisp set in terms of a pair of sets which is the lower and upper approximation of the original set. Let U denote the set of objects called universe and let R be an equivalence relation on U. Then (U, R) is called an approximation space. For  $u, v \in U \& (u, v) \in R$ , u and v belong to the same equivalence class it is denoted by U/R and we say that they are indistinguishable. The relation R is called an indiscernibility relation. Let  $[x]_R$  denote an equivalence class of R containing element x then Lower approximation  $\underline{R}(X)$  and upper approximation  $\overline{R}(X)$  for a subset  $X \subseteq U$  are defined by  $R(X) = \{x \in U/[x]_R \subset X\}$ 

$$\overline{R}(X) = \{x \in U/[x]_R \cap X \neq \phi\}$$

Thus if an object  $x \in \underline{R}(X)$  then "x surely belongs to X". If  $x \in \overline{R}(X)$  then "x possibly belong to X". If  $\underline{R}(X) \& \overline{R}(X)$  are sets then  $R(X) = (R(X), \overline{R}(X))$  is called a rough set with respect to R.

# **3. ROUGH GROUP**

**Definition3.1:** Let (U, R) be an approximation space, \* be a binary operation defined on U. Let  $[x]_R$  denote an equivalence class of R containing element *x*. Let  $X \subseteq U$  and the rough set  $R(X) = (\underline{R}(X), \overline{R}(X))$  is called an *upper rough group* if it satisfies the following axioms.

A1) for all  $x, y \in X, x * y \in \overline{R}(X)$ 

A2) Association property hold in  $\overline{R}(X)$ 

A3) there exist  $e \in \overline{R}(X)$  such that for all  $x \in X$ . x \* e = e \* x = x, e is called the rough identity element of X.

A4) for all  $x \in X$  there exist  $y \in \overline{R}(X)$ . x \* y = y \* x = e, y is called the rough inverse element of x in X.

**Definition 3.2:** Let (U, R) be an approximation space, \* be a binary operation defined on U. Let  $[x]_R$  denote an equivalence class of R containing element *x*. Let  $X \subseteq U$  and  $R(X) = (\underline{R}(X), \overline{R}(X))$  is called *a rough group* of U, if  $\underline{R}(X)$  and  $\overline{R}(X)$  are groups.

#### Jesmalar .L; International Journal of Advance Research, Ideas and Innovations in Technology.

#### **Observation 3.3**

- Let n  $(R_{\mu}(G))$  be the number of upper rough group and n(R (G)) be the number of rough group.
- The number of upper rough group of  $(Z_n, +_n)$  under the congruence relation (congruence modulo n/2). If n (n>3) is (a) even then
- $n(R_u(Z_4)) = 2^2 + 4 + 2.$ (i)
- $n(R_u(Z_6)) = 2^3 + 3(2^2) + 6 + 2.$ (ii)
- $n(R_u(Z_8)) = 2^4 + 4(2^3) + 6(2^2) + 8 + 2.$ (iii)
- $n(R_u(Z_{10})) = 2^5 + 5(2^4) + 8(2^3) + 8(2^2) + 10 + 2$ (iv)
- $n(R_u (Z_{12})) = 2^6 + 6(2^5) + 10(2^4) + 10(2^3) + 10(2^2) + 12 + 2$

In general  

$$n(R_u(Z_n)) = 2^{\frac{n}{2}} + \frac{n}{2} \left(2^{\frac{n}{2}-1}\right) + (n-2) \left(2^{\frac{n}{2}-2}\right) + \dots + (n-2)(2^2) + n + 2$$

- The number of upper rough group of  $(Z_n,+_n)$  under the congruence relation(congruence modulo $\frac{n-1}{2}$ ) If n(n>3) is odd then (b)
- (i)  $n(R_u (Z_5)) = 6(2) + 5 + 3.$

(v)

- (ii)  $n(R_{\nu}(Z_7)) = 6(2^2) + 12(2) + 2^2 + 7 + 3$
- (iii)
- (iv)
- $\begin{array}{l} n(R_u\ (Z_9))=6(2^3)+18(2^2)+18(2)+2^3+2^2+9+3\\ n(R_u\ (Z_{11}))=6(2^4)+24(2^3)+24(2^2)+24(2)+2^4+2^3+2^2+11+3\\ n(R_u\ (Z_{13}))=6(2^5)+30(2^4)+30(2^3)+30(2^2)+30(2)+2^5+2^4+2^3+2^2+13+3\\ \end{array}$ (v) In general  $n(R_u(\mathbb{Z}_n)) = 6(2^{\frac{n-3}{2}}) + 6\left(\frac{n-3}{2}\right)\left(2^{\frac{n-5}{2}}\right) + \dots + 6\left(\frac{n-3}{2}\right)\left(2^{\frac{n-(n-2)}{2}}\right) + 2^{\frac{n-3}{2}} + \dots + \left(\left(2^{\frac{n-(n-4)}{2}}\right)\right) + n + 3$

#### **Observation 3.4**

- (a) The number of rough group of  $(Z_n, +_n)$  under the congruence relation (congruence modulo n/2) If n(n>3) is even then
- $n(R(Z_4)=.\binom{2}{1})$ (i)
- $n(R(Z_6) = {\binom{2}{1}}^2$ (ii)
- $n(R(Z_8)) = {\binom{2}{1}} + {\binom{2}{1}}^3$  $n(R(Z_{10})) = ({\binom{2}{1}})^4$ (iii)
- (iv)
- $n(R(Z_{12})) = {\binom{2}{1}} + {\binom{2}{1}}^5$ (v)

In general 
$$n(R(Z_n)) = \begin{cases} \binom{2}{1} + \binom{2}{1}^{\frac{n}{2}-1} & \text{if } \frac{n}{2} \text{ is even} \\ \binom{2}{1}^{\frac{n}{2}-1} & \text{if } \frac{n}{2} \text{ is odd} \end{cases}$$

The number of rough group of  $(Z_n, +_n)$  under the congruence relation (congruence modulo $\frac{n-1}{2}$ ) If n(n>3) is odd then (b)

 $n(R(Z_n) = 0$ 

Theorem 3.5: Let (U, R) be an approximation space, \* be a binary operation defined on U. And U be a group with respect to \*. If  $\overline{R}(X)$  is a sub group then X is an upper rough subgroup. Proof

Since  $\overline{R}(X)$  is a subgroup and  $X \subseteq \overline{R}(X)$  for all  $x, y \in X, x * y \in \overline{R}(X)$ . Association property hold's in $\overline{R}(X)$ . And there exist (i)  $e \in \overline{R}(X)$  such that for all  $x \in X$ . x \* e = e \* x = x, e is called the rough identity element of X. for all  $x \in X$  there exist  $y \in \overline{R}(X)$  $\overline{R}(X), x * y = y * x = e, y$  is called the rough inverse element of x in X. Thus X is an upper rough subgroup of U. Remark 3.6

The converse of the above theorem need not be true.

for example.

Let U be the set of all permutation of  $S_4$  and \*be the multiplication operation of permutation. A classification of *U* is  $U/R = \{E_1, E_2, E_3, E_4\}$ , where

 $= \{(1), (12), (13), (14), (23), (24), (34)\},\$  $E_1$  $E_2$  $= \{(123), (132), (124), (142), (134), (143), (234), (243)\},\$ E3  $= \{(1234), (1243), (1324), (1342), (1423), (1432)\},\$  $E_4$  $= \{(12)(34), (13)(24), (14)(23)\},\$ Let  $X = \{(12), (123), (132)\}$ , then  $\overline{R}(X)$  $= E_1 \cup E_2$ 

Jesmalar .L; International Journal of Advance Research, Ideas and Innovations in Technology.

(i)  $\forall x, y \in X, x * y \in \overline{R}(X)$ 

- (ii) (12) \* (12) = (1)  $\in \overline{R}(X);$
- (iii) Association property holds in  $\overline{R}(X)$ ;

Thus we have X is an upper rough subgroup.

(iv)  $(12)^{-1} = (12) \in \overline{R}(X), (123)^{-1} = (132) \in \overline{R}(X),$ 

$$(132)^{-1} = (123) \in \overline{R}(X).$$

Here  $(123),(234) \in \overline{R}(X)$ , but $(123) * (234) = (13)(24) \notin \overline{R}(X)$ . Hence  $\overline{R}(X)$  is not a sub group.

## 4. HOMOMORPHISM AND ISOMORPHISM OF ROUGH GROUP

Let (U, R), (U'R') be two approximation spaces. Let U and U' be the groups with respect to the binary operations  $*, \overline{*}$  respectively. Let  $X \subseteq U, Y \subseteq U'$  such that  $\overline{R}(X), \overline{R}(X')$  be two sub groups of U, U' respectively. If  $f: \overline{R}(X) \to \overline{R}(X')$  is a homomorphism, then we say that  $f: X \to X'$  is an *upper rough homomorphism* such that for all  $x, y \in \overline{R}(X)$ , f(x \* y) = f(x) \* f(y).

**Theorem 4.1:** Let (U, R), (U', R') be two approximation spaces, Let *U* and *U*'be the groups with respect to the binary operations  $*, \overline{*}$  respectively. Let  $X \subseteq U, Y \subseteq U'$  such that  $\overline{R}(X), \overline{R}(X')$  are two sub group of *U*, *U*'respectively. If *f*:  $\overline{R}(X) \to \overline{R}(X')$  is a homomorphism,  $f: X \to X'$  is an upper rough homomorphism, then the following properties hold.

(i) 
$$f(e) = e^{-e^{i\theta}}$$

(ii)  $f(a^{-1}) = [f(a)]^{-1}$ 

## Proof

(i) Let  $a \in X$  then  $a \in \overline{R}(X)$  there exist  $e \in \overline{R}(X)$  such that a \* e = e \* a = e then f(a \* e) = f(e \* a) = f(a)f(a) \* f(e) = f(e) \* f(a) = f(a)

Hence f(e) = e',  $e' \in \overline{R}(X')$ (ii) Let  $a \in X$  then  $a^{-1} \in \overline{R}(X)$  such that  $a * a^{-1} = a^{-1} * a = e$   $f(a * a^{-1}) = f(a^{-1} * a) = f(e)$   $f(a) = f(a^{-1}) = f(a^{-1}) = f(a) = f(e) = e'$ Hence  $f(a^{-1}) = [f(a)]^{-1}$ 

**Definition 4.2:** Let(U, R),(U'R') be two approximation spaces. Let U and U' be the groups with respect to the binary operations  $*, \overline{*}$  respectively.  $X \subseteq U, X' \subseteq U'$  such that  $\overline{R}(X), \overline{R}(X')$  are sub groups of U, U' respectively. If  $f: \overline{R}(X) \to \overline{R}(X')$  is a isomorphism, then we say that  $f: X \to X'$  is an *upper rough Isomorphism*.

**Theorem 4.3:** Let (U, R), (U', R') be two approximations spaces, where U and U' are groups with respect to the binary operations  $*, \overline{*}$  respectively. Let  $X \subseteq U, X' \subseteq U'$  such that  $\overline{R}(X), \overline{R}(X')$  are sub groups of U, U' respectively. If  $f: \overline{R}(X) \to \overline{R}(X')$  is an isomorphism, then

(i) 
$$f\left(\overline{R}(X)\right) = \overline{R}(f(X))$$
  
(ii)  $f^{-1}\left(\overline{R}(X)\right) = \overline{R}(f^{-1}(X))$ 

## Proof

(i)  $f\left(\overline{R}(X)\right) = \overline{R}(f(X))$ Let  $x \in f(\overline{R}(X))$  $\Longrightarrow f^{-1}(x) \in \overline{R}(X),$  $\Rightarrow [f^{-1}(x)]_R \cap X \neq \phi$  $\Rightarrow f([f^{-1}(x)]_R \cap X) \neq \phi$  $\Rightarrow f([f^{-1}(x)]_R) \cap f(X) \neq \phi$  $\Rightarrow [x]_R \cap f(x) \neq \phi$ [since f is an isomorphism]  $\Rightarrow x \in \overline{R}(f(X))$  $f\left(\overline{R}(X)\right) \subseteq \overline{R}(f(X))$ Let  $x \in \overline{R}(f(X))$  $\Rightarrow x \in [f(x)]_R \cap f(X)$  $\Rightarrow x \in f(X)$  $\Rightarrow x \in f\left(\overline{R}(X)\right) [\text{since } X \subseteq \overline{R}(X)] \Rightarrow f(X) \subseteq f(\overline{R}(X))]$  $\overline{R}(f(X)) \subseteq f\left(\overline{R}(X)\right)$ Hence  $f(\overline{R}(X)) = \overline{R}(f(X))$ 

$$\begin{aligned} \text{(ii)} f^{-1}\left(\overline{R}(X)\right) &= \overline{R}(f^{-1}(X)) \\ \text{Let } x \in f^{-1}\left(\overline{R}(X)\right) \\ &\Rightarrow f(x) \in \overline{R}(X), \\ &\Rightarrow [f(x)]_R \cap X \neq \phi \\ &\Rightarrow f^{-1}([f(x)]_R \cap X) \neq \phi \\ &\Rightarrow f^{-1}([f(x)]_R) \cap f^{-1}(X) \neq \phi \\ &\Rightarrow [x]_R \cap f^{-1}(x) \neq \phi \qquad \text{[since f is an isomorphism]} \\ &\Rightarrow x \in \overline{R}(f^{-1}(X)) \\ f^{-1}\left(\overline{R}(X)\right) \subseteq \overline{R}(f^{-1}(X)) \\ \text{Let } x \in \overline{R}(f^{-1}(X)) \\ &\Rightarrow x \in f^{-1}(\overline{R}) \\ &\Rightarrow x \in f^{-1}(\overline{R}) \text{[Since} X \subseteq \overline{R}(X) \text{]} \Rightarrow f^{-1}(X) \subseteq f^{-1}(\overline{R}(X)) \text{]} \\ &\overrightarrow{R}(f^{-1}(X)) \subseteq f^{-1}\left(\overline{R}(X)\right) \\ &\text{Hence } f^{-1}\left(\overline{R}(X)\right) = \overline{R}(f^{-1}(X)) \end{aligned}$$

**Theorem 4.4:** Let (U, R), (U', R') be two approximation spaces, Let *U* and *U*' be the groups with respect to the binary operations  $*, \overline{*}$  respectively. Let  $X \subseteq U, X' \subseteq U'$  such that  $\overline{R}(X), \overline{R}(X')$  be two sub group of *U*, *U*' respectively. If *f*:  $\overline{R}(X) \to \overline{R}(X')$  is an isomorphism,  $f : X \to X'$  is an upper rough isomorphism, then the following properties hold.

- (i) fIf  $X_1$  is an upper rough sub group of X then  $f(X_1)$  is an upper rough subgroup of X'
- (ii) If  $X_1$  is an upper rough normal in X then  $f(X_1)$  is an upper rough normal in f(X)
- (iii) If  $X_1'$  is an upper rough sub group of X' then  $f^{-1}(X_1')$  is an upper rough subgroup of X
- (iv) If  $X_1'$  is an upper rough normal in X' then  $f^{-1}(X_1')$  is an upper rough normal in X

# Proof

(i) If  $X_1$  is an upper rough sub group of X Since  $X_1$  is non empty,  $f(X_1)$  is also non empty. Now let  $x, y \in f(X_1)$  then  $x, y \in f(\overline{R}(X_1))$ , x = f(a), y = f(b) where  $a, b \in \overline{R}(X_1)$ Therefore  $x = y^{-1} = f(a) = [f(b)]^{-1} = f(a) = f(a) = f(a + b^{-1})$ Since  $X_1$  is an upper rough sub group of X,  $a * b^{-1} \in \overline{R}(X_1)$ Therefore  $x = y^{-1} = f(a * b^{-1}) \in f(\overline{R}(X_1))$ Therefore  $f(\overline{R}(X_1))$  is a subgroup of X' By Theorem 4.3  $f(\overline{R}(X)) = \overline{R}(f(X))$  $\overline{R}(f(X_1))$  is a subgroup of X' By Theorem 3.5  $f(X_1)$  is an upper rough subgroup of X' (ii) Let  $X_1$  is an upper rough normal in X Let  $x \in f(X_1)$  and  $y \in f(X)$ claim $y * x * y^{-1} \in \overline{R}(f(X_1))$ Now, x = f(a), y = f(b) where  $a \in X_1 \& b \in X$ Since  $X_1$  is an upper rough normal in X,  $\overline{R}(X_1)$  is normal in,  $b * a * b^{-1} \in \overline{R}(X_1)$ Therefore  $f(b * a * b^{-1}) \in f(\overline{R}(X_1))$  $f(b) = \overline{f(a)} = f(b^{-1}) \in f(\overline{R}(X_1))$ Therefore  $y = x = y^{-1} \in f(\overline{R}(X_1))$  $f(\overline{R}(X_1))$  is normal in f(X)By Theorem 4.3  $\overline{R}(f(X_1))$  is normal in f(X) $\Rightarrow f(X_1)$  is an upper normal in f(X)(iii) Let  $X_1'$  be an upper rough sub group of X'Since  $(e) = e' \in \overline{R}(X_1')$ ,  $e \in f^{-1}(\overline{R}(X_1'))$ hence  $f^{-1}(\overline{R}(X_1')) \neq \varphi$ Let  $a, b \in f^{-1}(\overline{R}(X_1'))$  then  $f(a), f(b) \in \overline{R}(X_1')$  $\Rightarrow f(a) = [f(b)]^{-1} \in \overline{R}(X_1')$ 

 $\Rightarrow f(a * b^{-1}) \in \overline{R}(X_1')$   $\Rightarrow a * b^{-1} \in f^{-1}(\overline{R}(X_1'))$   $f^{-1}(\overline{R}(X_1')) \text{ is a subgroup of } X$   $\Rightarrow \overline{R}(f^{-1}(X_1') \text{ is sub group of } X.$   $\Rightarrow f^{-1}(X_1') \text{ is an upper rough subgroup of } X$ Hence  $f^{-1}(X_1'')$  is an upper rough subgroup of X(iv) Let  $X_1'$  is an upper rough normal in X'

Let  $\in f^{-1}(X_1')$  and  $a \in X$ Then  $f(x) \in X_1'$  and  $f(a) \in f(X)$ To prove  $a * x * a^{-1} \in f^{-1}(X_1')$ 

Since  $X_1'$  is an upper rough normal in X',  $\overline{R}(X_1')$  is normal in X'

$$\Rightarrow f(a) \stackrel{\overline{*}}{=} f(x) \stackrel{\overline{*}}{=} [f(a)]^{-1} \in \overline{R}(X_1') \Rightarrow f(a * x * a^{-1}) \in (\overline{R}(X_1)) \Rightarrow a * x * a^{-1} \in f^{-1}(\overline{R}(X_1'))$$

 $f^{-1}(\overline{R}(X_1'))$  is normal in X  $\Rightarrow \overline{R}(f^{-1}(X_1'))$  is normal in X [by theorem 4.3]  $f^{-1}(X_1')$  is an upper rough normal in X.

**Theorem 4.5:** An upper rough isomorphism is an equivalence relation **Proof** 

Let  $\overline{R}(X),\overline{R}(X')$  be two sub groups of U and U' respectively, then by theorem 3.5, X and X' be two upper rough sub groups of U and U' respectively.

For any group  $\overline{R}(X)$ ,  $i_x = \overline{R}(X) \to \overline{R}(X')$  is an isomorphism clearly the identity map  $i_x = X \to X'$  is an upper rough isomorphism,

hence  $X \cong_r X$ 

Therefore the relation is reflexive.

Let  $\overline{R}(X),\overline{R}(X')$  be two sub group of U, U' respectively, then X, X' be two upper rough sub group of U, U' respectively

Now let  $X \cong_r X'$  and let  $f : \overline{R}(X) \to \overline{R}(X')$  be an isomorphism,  $\Rightarrow f: \overline{R}(X) \rightarrow \overline{R}(X')$  is a bijection.  $\Rightarrow f^{-1}: \overline{R}(X') \rightarrow \overline{R}(X)$  is also a bijection Let  $x', y' \in \overline{R}(X')$ Since f is onto there exists  $x \in X, y \in X$ such that x' = f(x) and y' = f(y)  $\Rightarrow x = f^{-1}(x')$  and  $y = f^{-1}(y')$ Therefore f(xy) = f(x)f(y) = x'y'Therefore  $f^{-1}(x'y') = xy = f^{-1}(x')f^{-1}(y')$ Therefore  $f^{-1}$ :  $\overline{R}(X') \to \overline{R}(X)$  is a homomorphism Therefore  $f^{-1}$ :  $\overline{R}(X') \to \overline{R}(X)$  is an isomorphism Hence  $f^{-1}: X' \to X$  is an upper  $X' \cong_R X$ rough isomorphism Hence and The relation sis symmetric Now let  $X \cong_r X'$  and  $X' \cong_r X''$  and let  $f : \overline{R}(X) \to \overline{R}(X')$ be an isomorphism,  $h: \overline{R}(X') \to \overline{R}(X'')$  is an isomorphism.  $hof: \overline{R}(X) \to \overline{R}(X')$  is also rough bijection Now let  $x, y \in X$  then (hof)(xy) = h[f(xy)]= h[f(x)(y)]= h(f(x))h(f(y))= (hof)(x)(hof)(y)

 $hof: \overline{R}(X) \to \overline{R}(X'')$  is an isomorphism

Hence hof  $X \to X''$  is an upper rough isomorphism Thus  $X \cong_r X''$  and hence the relation is transitive. Therefore an upper rough isomorphism is an equivalence relation.

### REFERENCES

- 1. Biswas, R and Nanda, S. Rough groups and rough subrings, Bull. Polish Acad. Sci. Math, 42(1994), 251 154.
- Duaian Miao, Suing Han, Daoguo Li and Lijun Sun, "Rough group, Rough subgroup and their properties" Springer Volume 3641 of the series of Lecture Notes in Computer Science pp104-113
- 3. Herstein, I.N., Topics in Algebra, 2nd edition, Wiley, New York, 1975.
- 4. Kuroki, N. Rough ideals in semigroups, Infor. Sci. 100(1997), 136 163. DOI: 10.1016/S0020-0255(96)00274-5

## Jesmalar .L; International Journal of Advance Research, Ideas and Innovations in Technology.

- 5. Pawlak, Z. Rough set theory and its applications, Journal of telecommunications and information technology, 3/2002.
- 6. Pawlak, Z. Rough sets, Theoretical aspects of reasoning about data, Kluwer Academic Publishers, Dordrecht, 1991.
- 7. Pawlak, Z. Rough sets, International Journal of computer and information sciences 11:341-356, 198211:341-356,1982
- S. Robinson Chellathurai & L. Jesmalar, Core in Rough Graph and Weighted Rough Graph, International Journal of Contemporary Mathematical Sciences, Vol. 11, 2016, no. 6, 251 - 265, ISSN (1312-7586), June-2016. Citation 0,165, H.Index: 12.
- 9. S. Robinson Chellathurai& L. Jesmalar, Rough lattice and Core in Rough data set lattice, Advances in Fuzzy Mathematics. ISSN 0973-533XVolume 11, Number 2 (2016), pp. 113-127, Sep2016.
- 10. S. Robinson Chellathurai & L. Jesmalar, Rough sets on Clustering, International Journal of Engineering and Computing, Volume.7, Issue No.3, PP 5032-5035, ISSN (2321-3361), March-2017. Impact factor:5.611